Division of Behavioral Health Services

Division of Compliance

150 North 18th Avenue, Suite 260 Phoenix, Arizona 85007 (602) 364-4670 (602) 364-4762 FAX

JANET NAPOLITANO, GOVERNOR SUSAN GERARD, DIRECTOR

TO: RBHA and TRBHA CEOs, Medical Directors and Clinical Leadership

FROM: Jennifer Vehonsky

Acting Division Chief, Bureau of Compliance

Internet: www.azdhs.gov

SUBJECT: POLICY CLARIFICATION: ARIZONA CROSSWALK FOR DC: 0-3R

AND ICD-9-CM

DATE: May 19, 2006

The purpose of this memorandum is to disseminate to RBHA and TRBHA staff, Medical Directors and Clinical staff a revised crosswalk of diagnosis codes for zero – three year olds (DC: 0-3R). The Arizona DC: 0-3R crosswalk serves strictly as a tool to assist the clinician in assigning the correct diagnostic code required for reimbursement when working with children under the age of three. The Arizona DC: 0-3R is included as a separate attachment.

This new edition Arizona DC: 0-3R includes the following changes:

- 1) "Gender Identity Disorder" was removed.
- 2) "Reactive Attachment Deprivation/Maltreatment Disorder of Infancy" was changed to "Deprivation/Maltreatment Disorder of Infancy."
- 3) "Traumatic Stress Disorder was renamed "Posttraumatic Stress Disorder."
- 4) "Regulatory Disorders" was renamed "Regulation Disorders of Sensory Processing."
- 5) Subtypes have been added to "Sleep Behavior Disorders" and "Eating Behavior Disorders."
- 6) Subtypes have been added to "Anxiety Disorder" for use in children over 2 years of age.
- 7) "Pervasive Developmental Disorder" can now be used for children over 2 years of age.

Arizona's Crosswalk for DC: 0-3R and ICD-9-CM

In 2005, ZERO TO THREE's *Diagnostic Classification of Mental Health and Developmental Disorders of Infancy and Early Childhood* (DC: 0-3) was updated and revised. This new edition, DC: 0-3R, includes the following changes:

- 1) "Gender Identity Disorder" was removed.
- 2) "Reactive Attachment Deprivation/Maltreatment Disorder of Infancy" was changed to "Deprivation/Maltreatment Disorder of Infancy."
- 3) "Traumatic Stress Disorder was renamed "Posttraumatic Stress Disorder."
- 4) "Regulatory Disorders" was renamed "Regulation Disorders of Sensory Processing."
- 5) Subtypes have been added to "Sleep Behavior Disorders" and "Eating Behavior Disorders."
- 6) Subtypes have been added to "Anxiety Disorder" for use in children over2 years of age.
- 7) "Pervasive Developmental Disorder" can now be used for children over 2 years of age.

Several states had previously developed crosswalks between the codes from DC: 0-3 and ICD-9-CM. With the changes noted above, however, a revised crosswalk is necessary.

The Arizona DC: 0-3R crosswalk serves strictly as a tool to assist the clinician in assigning the correct diagnostic code required for reimbursement when working with children under the age of three. If a DSM-IV-TR diagnosis best describes the primary presenting difficulty, it should be coded in Axis I and the corresponding ICD-9-CM code used. For example, if the diagnosis is Pica, Rumination Disorder, or another diagnosis not included in the DC: 0-3R, the clinician should list the appropriate DSM-IV-TR diagnosis and utilize the appropriate ICD-9-CM code.

Disclaimer: End dated diagnosis codes are not valid for current use. It is recommended that the current ICD-9-CM book be used in conjunction with this list to validate fourth and fifth digit requirements.

^{*} Per AHCCCS policy, Primary Diagnosis for billing purposes is restrict6ed to ICD-9-CM codes 290 through 316.99. Codes outside of this specified range cannot be used as Primary Diagnosis.

DC: 0-3 R Diagnosis		ICD-9-CM Diagnosis	
100	Post Traumatic Stress Disorder	308.0	Predominant disturbance of emotions ■ Anxiety, emotional crisis, or panic state as acute reaction to exceptional (gross) stress
		308.2	Predominant psychomotor disturbance Agitation states or stupor as acute reaction to exceptional (gross) stress
		308.3	Other acute reactions to stress Acute situational disturbance Acute stress disorder
		309.81	Posttraumatic stress disorder Chronic posttraumatic stress disorder Posttraumatic stress disorder NOS
150	Deprivation/Maltreatment Disorder	*995.50	Child abuse, unspecified
		*995.51 *995.52	Child neglect (nutritional)
		*995.52	Child neglect (nutritional) Child sexual abuse
		*995.54	Child physical abuse Battered baby or child syndrome
		*995.59	Other child abuse and neglect Multiple forms of abuse
200	Disorders of Affect		

^{*} Per AHCCCS policy, Primary Diagnosis for billing purposes is restrict6ed to ICD-9-CM codes 290 through 316.99. Codes outside of this specified range cannot be used as Primary Diagnosis.

210	Prolonged Bereavement/ Grief Reaction	309.0	Adjustment disorder with depressed mood Grief reaction
		309.1	Prolonged depressive reaction
220	Anxiety Disorders of Infancy and Early Childhood		
221	Separation Anxiety Disorder	309.21	Separation anxiety disorder
222	Specific Phobia	300.20	Phobia, unspecifiedAnxiety-hysteria NOSPhobia NOS
		300.29	 Other isolated or specific phobias Acrophobia, animal phobias, claustrophobia, or fear of crowds
223	Social Anxiety Disorder (Social Phobia)	300.23	Social phobia Fear of eating in public, speaking in public, washing in public
224	Generalized Anxiety Disorder	300.02	Generalized anxiety disorder
225	Anxiety Disorder NOS	300.00	Anxiety state, unspecified
230	Depression of Infancy and Early Childhood		

^{*} Per AHCCCS policy, Primary Diagnosis for billing purposes is restrict6ed to ICD-9-CM codes 290 through 316.99. Codes outside of this specified range cannot be used as Primary Diagnosis.

231	Type I: Major Depression	296.2x	Major depressive disorder, single episode
			Fifth digit must be inserted:
232	Type II: Depressive Disorder NOS	311	 Depressive disorder, not elsewhere classified Depressive disorder NOS, depressive state NOS, depression NOS
240	Mixed Disorder of Emotional Expressiveness	313.9	Unspecified emotional disturbance of childhood or Adolescence Mental disorder of infancy, childhood, or adolescence NOS
300	Adjustment Disorder	309.0	Adjustment disorder with depressed mood Grief reaction
		309.24	Adjustment disorder with anxiety
		309.28	Adjustment disorder with mixed anxiety and depressed mood
		309.3	Adjustment disorder with disturbance of conduct Conduct disturbance or destructiveness as adjustment reaction
		309.4	Adjustment disorder with mixed disturbance of emotions and conduct

^{*} Per AHCCCS policy, Primary Diagnosis for billing purposes is restrict6ed to ICD-9-CM codes 290 through 316.99. Codes outside of this specified range cannot be used as Primary Diagnosis.

		308.82	Adjustment reaction with physical symptoms
		309.83	Adjustment reaction with withdrawal Elective mutism as adjustment reaction Hospitalism (in children) NOS
		309.89	Adjustment Reaction, Other
		309.9	 Unspecified adjustment reaction Adaptation reaction NOS Adjustment reaction NOS
400	Regulation Disorders of Sensory Processing		
410	Hypersensitive		
411	Type A: Fearful/Cautious	313.0	Overanxious disorder Anxiety and fearfulness or overanxious disorder of childhood and adolescence
412	Type B: Negative Defiant	313.81	Oppositional defiant disorder
420	Hyposensitive/Underresponsive	313.21	Shyness disorder of childhood Sensitivity reaction of childhood or adolescence
		313.22	Introverted disorder of childhood Social withdrawal or withdrawal reaction of childhood or adolescence
430	Sensory Stimulation- Seeking/Impulsive	314.01	Attention deficit disorder with hyperactivity

^{*} Per AHCCCS policy, Primary Diagnosis for billing purposes is restrict6ed to ICD-9-CM codes 290 through 316.99. Codes outside of this specified range cannot be used as Primary Diagnosis.

		314.1	 Hyperkinesis with developmental delay Developmental disorder of hyperkinesis Use additional code to identify any associated neurological disorder
			Unspecified hyperkinetic syndrome
			 Hyperkinetic reaction of childhood or adolescence NOS
500	Class Bahavias Biografias		Hyperkinetic syndrome NOS
500	Sleep Behavior Disorder	207.40	Newsyspia alon discader unemodified
510	Sleep-Onset Disorder (Protodyssomnia)	307.40	Nonorganic sleep disorder, unspecified
		307.41	Transient disorder of initiating or maintaining sleep
520	Night-Waking Disorder		 Adjustment insomnia
	(Protodyssomnia)		 Hyposomnia, insomnia, or sleeplessness associated with intermittent emotional reactions or conflicts
		307.42	 Persistent disorder of initiating or maintaining sleep Hyposomnia, insomnia, or sleeplessness associated with: anxiety, conditioned arousal, depression (major)(minor), psychosis Idiopathic insomnia Paradoxical insomnia Primary insomnia Psychophisioogical insomnia
		307.45	Circadian rhythm sleep disorder of nonorganic origin
		307.46	Sleep arousal disorder

^{*} Per AHCCCS policy, Primary Diagnosis for billing purposes is restrict6ed to ICD-9-CM codes 290 through 316.99. Codes outside of this specified range cannot be used as Primary Diagnosis.

			 Night terror disorder, night terrors, sleep terror disorder, sleepwalking, somnambulism
600	Feeding Behavior Disorder		
601	Feeding Disorder of State Regulation	307.50	Eating Disorder, unspecified ■ Eating disorder, NOS
602	Feeding Disorder of Caregiver- Infant Reciprocity	307.59	Other feeding disorder of infancy or early childhood of nonorganic origin
603	Infantile Anorexia		 infantile feeding disturbances or loss of appetite of nonorganic origin
604	Sensory Food Aversions	*783.3	Feeding difficulties and mismanagement
605	Feeding Disorder Associated with Concurrent Medical Condition	*779.3	Regurgitation, slow feeding, vomiting in newborn
606	Feeding Disorder Associated with Insults to the Gastrointestinal Tract		
700	Disorders of Relating and Communicating		
710	Multisystem Developmental Disorder (MSDD) for children under the age of two years; Pervasive Developmental Disorders can be utilized for children over the age of two if identified	299.0x	Autistic disorder

^{*} Per AHCCCS policy, Primary Diagnosis for billing purposes is restrict6ed to ICD-9-CM codes 290 through 316.99. Codes outside of this specified range cannot be used as Primary Diagnosis.

299.9x	 Unspecified pervasive developmental disorder Child psychosis NOS Schizophrenia, childhood type, NOS Schizophrenia syndrome of childhood NOS
	Fifth digit must be inserted: O current or active state 1 residual state

^{*} Per AHCCCS policy, Primary Diagnosis for billing purposes is restrict6ed to ICD-9-CM codes 290 through 316.99. Codes outside of this specified range cannot be used as Primary Diagnosis.